

Detection of Knots and Jets in IRAS 06061+2151

B.G. Anandarao¹ A. Chakraborty² D.K. Ojha³ and L. Testi⁴

¹ Physical Research Laboratory, Ahmedabad-390009, India

² Department of Astronomy & Astrophysics, 525 Davey Laboratory, Pennsylvania State University, University Park, PA 16802, USA

³ Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai-400005, India

⁴ Osservatorio Astrofisico di Arcetri, INAF, Largo E. Fermi 5, 50125, Firenze, Italy

Abstract. We report detection of a Young Stellar Object with an evidence for an outflow in the form of knots in the molecular hydrogen emission line ($2.121\mu\text{m}$) towards the massive star forming region IRAS 06061+2151. Near-infrared images reveal IRAS 06061+2151 to be a cluster of at least five sources, four of which seem to be early B type young stellar objects, in a region of 12 arcsecs surrounded by a nebulosity. The presence of the knots that are probably similar to the HH objects in the optical wavelengths, suggests emerging jets from one of the cluster members. These jets appear to excite a pair of knot-like objects (Knot-NW and Knot-SE) and extend over a projected size of 0.5pc. The driving source for the jets is traced back to a member of the cluster whose position in the H-Ks/J-H color-color diagram indicates that it is a Class I type pre-mainsequence star. We also obtained K band spectra of the brightest source in the cluster and of the nearby nebular matter. The spectra show molecular hydrogen emission lines but do not show Br γ line ($2.167\mu\text{m}$). These spectra suggest that the excitation of the molecular hydrogen lines is probably due to a mild shock.

Key words. stars: pre-mainsequence, ISM: jets and outflows; stars: formation

1. Introduction

IRAS 06061+2151 is one of the most luminous IRAS sources believed to be associated with the galactic massive star forming region Gemini OB1 which is at a distance of 2 kpc (Carpenter, Snell & Schloerb 1995a). The object figures in the radio survey of massive star forming regions done by Kurtz, Churchwell & Wood (1994; hereafter KCW94). The CO survey of Shepherd & Churchwell (1996) categorized this object as having a moderate velocity gas flow (full width = 20.8 km/s and fwhm = 4 km/s). This Ultra Compact HII(UCHII) region was previously studied by Carpenter, Snell & Schloerb (1995a) and very recently in a near-infrared (NIR) survey on UCHII regions by Hanson, Luhman & Rieke (2002; hereafter HLR02). The extensive CS survey and NIR observations by Carpenter, Snell & Schloerb (1995a,b) of 58 IRAS sources in the Gemini OB1 molecular cloud complex suggested that more luminous IRAS sources tend to be associated with more massive cores. Based on their study, these authors proposed a qualitative model for the formation of massive dense cores, primarily through the external compression that sweeps shells of molecular gas.

The mass outflows in the form of jets from young stellar objects (YSOs) are the important signatures of star formation in molecular clouds (Hartmann 1998 and references therein). The jets in turn cause Herbig-Haro (HH) objects seen in optical emission lines notably [SII] and H α by shocking/interacting with the ambient interstellar medium. Such manifestations are better revealed in the NIR due to the lesser extinction that it suffers compared to the visible region. In the recent past several discoveries have been made of the molecular hydrogen outflows; both in situations where there exist optical counterparts in the form of HH objects (eg., Schwartz et al 1988; Davis, Eisloffel & Ray, 1994; Eisloffel, Smith & Davis 2000) and where there are no optical counterparts (eg., Herbst, Beckwith & Robberto 1997). However, the outflows detected in the case of massive proto-stars are rare due to the fact that the outflows are short-lived compared to those in the low mass proto-stars, though it is believed that stars of all masses might undergo these phenomena.

Therefore, luminous IRAS sources such as the IRAS 06061+2151 are the appropriate candidates for further study in the NIR. Our motivation is to look for new outflows associated with massive protostars and to determine their evolutionary stages (such as Class I, II and III), using color-color diagrams (Lada & Adams 1992; Lada, Young & Greene 1993; Gomez, Kenyon & Hartmann 1994).

In this paper, we report detection of infrared counterparts of HH objects presumably caused by jets from a class I type source towards IRAS 06061+2151. We study the nature of the cluster of sources using JHK photometry, narrowband images in molecular hydrogen ($2.12\mu\text{m}$) and $\text{Br}\gamma$ lines and medium resolution K band spectroscopy. The data were obtained by new observations as well as from the 2MASS archive. In section 2 we present the details of observations and data reduction procedures, section 3 deals with results and discussion and we summarize our conclusions in section 4.

2. Observations and Data Reduction

2.1. Observations from Mt. Abu

J($1.253\mu\text{m}/0.296\mu\text{m}$), H($1.659/0.288$), K'($2.120/0.360$) photometric observations of IRAS 06061+2151 were done from the 1.2 meter PRL Infrared Telescope, Mt. Abu, India, using a 256×256 HgCdTe array (NICMOS-3, made by Infrared Laboratories, Arizona, USA). The instrument is described in Anandarao (1998) and Nandakumar (1999). The plate scale is $0.5''/\text{pixel}$ (corresponding to a field of view of $2'$). The night of observation (26 Jan 2001) was photometric with a seeing of $0.9\text{--}1.0''$.

Individual object frames were of 30s of integration in J and H bands and 3s in K' band. The data reduction was done using IRAF software tasks. All the images were subjected to standard pipeline procedures like dark and sky subtraction and flat-fielding. The images were then co-added to obtain a final image in each band (J, H and K') averaged over a total integration time of 300s. Further, in each band two such images were obtained. The images were then filtered using a median filter of 3×3 pixels for removing noise at sky-level, thus making them suitable for photometric analysis. We used DAOPHOT (Stetson 1987) task and followed the photometric procedure described by Chakraborty et al. (2000). We estimate the overall photometric error to be ± 0.08 mag. The faintest stars recorded were found to be of 16.66, 15.30 and 14.10 magnitudes in J, H and K' bands respectively.

2.2. 2MASS data

We extracted sources from the 2MASS All-Sky Point Source Catalogue (PSC) which were within $2'$ radius around the IRAS source. Thus this region excludes the cluster G188.80+01.07 (see Bica, Dutra & Barbuy 2003). The data were downloaded from the 2MASS Homepage available for the Astronomical community. The 2MASS point source catalogue consists of J($1.24\mu\text{m}/0.33\mu\text{m}$), H($1.66/0.39$) and Ks($2.16/0.42$) magnitudes of stars. The PSC detection limits for a $S/N=10$ are $J=15.8$, $H=15.1$ and $Ks=14.3$.

The 2MASS data are used to produce the color-color diagram to study the nature of sources. We use the Mt. Abu photometry for the five bright cluster candidate YSOs towards IRAS 06061+2151 (see Sec. 3.1). It is to

be noted that the Mt. Abu magnitudes match reasonably well with the 2MASS data as shown in Table 1 (except for Stars 1 and 5), inspite of the differences in the passbands.

2.3. Observations from TNG, La Palma

We obtained spectra ($R=1000$, 1.4 to $2.4\mu\text{m}$) of one of the bright sources (Star no. 1) in the cluster of IRAS 06061+2151 using the Near Infrared Camera Spectrometer (NICS) attached to the 3.58 m Telescopio Nazionale Galileo (TNG) at La Palma. NICS has a HgCdTe Hawaii 1024x1024 array and it was used under LF configuration with a plate scale of $0.25''$ (for instrument details, see Baffa et al. 2001). The medium resolution spectra were obtained using the HK grism with a slit width of $0.5''$. The Position Angle (PA) of the slit was 270 degrees i.e., in the East-West direction. Two dithered exposures of 60 secs each were obtained on the Star no. 1. A hot star (A0) spectra were also taken for telluric calibration. Following a similar procedure, spectra were obtained in a region $5''$ west of Star no. 1, in order to sample the nebular regions.

Apart from the spectra, we also obtained images of the source in two narrowband filters centered around $2.12\mu\text{m}$ for the H_2 $v=1-0$ S(1) line and $2.17\mu\text{m}$ for the $\text{Br}\gamma$ line. A total of five 60 seconds dithered exposures (dithered by more than 3 arcmins) were obtained in both the narrowband filters. The seeing was $0.8\text{--}0.9$ arcsec during observations.

The TNG images were analysed in a similar process as those from Mt. Abu. The spectra were reduced using standard IRAF tasks like APALL, TELLURIC, and normalized with respect to the respective continua. The wavelength calibration was done using the sky OH lines using the list of lines available with IRAF. We present here the K band spectra for which the signal to noise ratio is better than 5. This spectral region contains some of the most important features that we are looking for, such as the $\text{Br}\gamma$ and H_2 vibrational lines.

3. Results and Discussion

3.1. Photometry

Fig 1 shows the K' band image of the central region of the object taken at Mt. Abu. The five prominent stellar sources (S1,2,3,4 and 5) are marked on the figure and their magnitudes are given in Table 1. The data for the S4 (all three bands) and S5 (only in K' band) are from the Mt. Abu observations. S4 is detected in H and Ks bands but not in J band; while S5 is detected only in Ks band in 2MASS. The Mt. Abu K' magnitude of S5 is 12.9 while in J and H bands the source was below the detection limit. HLR02 obtained an H band image of the central region of this object which does not show S5; thus lending support our result. It appears then that this source has a rather steep spectral energy distribution in the NIR and beyond. Hence the importance of the NIR study at high angular resolu-

tion (atleast seeing-limited) of the massive star forming regions.

The IRAS flux densities for 06061+2151 are 12.0, 144.8, 865.6 and 1130 Jy in the 12, 25, 60 and 100 μm bands respectively. From these values, Carpenter, Snell & Schloerb (1995b) estimated the total far-infrared (FIR) luminosity to be $4000 L_{\odot}$. Based on the total FIR luminosity, Ghosh et al. (2000) suggested that the core may contain a single B0-B0.5 type hot star, though they could not resolve the cluster due to the large field of view (1.5 arcmin). Our NIR image (see Figure 1), however, reveals that IRAS 06061+2151 consists of a cluster of at least five stars.

Figure 2 shows J-H/H-Ks color-color diagram of sources from the 2MASS data and Mt. Abu observations (S1-S4). The 2MASS-PSC data covers a diameter of 4 arcmins around the co-ordinate of the IRAS source. The total number of sources extracted were 57, after removing all the "null" entries. The solid curve is the locus of points corresponding to unreddened main sequence stars and giants (Bessell & Brett 1988). The two parallel lines (dashed) are the reddening vectors drawn upto the visual extinction of $A_V = 30$; together these parallel lines show the region in which the main-sequence stars occur at various extinctions (Hunter et al. 1995). Sources on the right side of the reddening strip (~ 17) are primarily stars with intrinsic color excess which include the S1-S4 stars of the cluster. We note here that we allowed an uncertainty of 0.1 mag in H-K color on either side of the strip to identify the left and right side of the reddening strip. Also shown in the figure is the locus for the CTTS (from Meyer, Calvet & Hillenbrand 1997) and the corresponding reddening vector.

The stars on the right side are mostly Young Stellar Objects (YSOs) (Lada & Adams 1992; Lada, Young & Greene 1993; Gomez, Kenyon & Hartmann 1994). In general, sources whose H-K color is ≥ 1.5 and J-H ≥ 1.0 can be considered as Class I type YSOs or protostars in the early stages of their evolution (Strom, Margulis & Strom 1989; Kenyon et al. 1993; Lada & Adams 1992). From Fig 2, we find that out of the total of 17 stars falling clearly in the region on right side of the reddening vector, 3-4 are clearly beyond the CTTS locus to its right. These stars that include S4 and maybe S2 can be classified as possible Class I candidates, if we account the uncertainty of 0.1 mag in colors. The rest numbering about 13-14 stars can be classified as possible Class II - III. Thus, the region around IRAS 06061+2151 shows a concentration of YSOs. Indeed, NIR studies by Carpenter, Snell & Schloerb (1995b) predicted a similar trend towards IRAS 06061+2151.

Figure 3 shows H-Ks/Ks color-magnitude diagram. Like in figures 1 and 2 the stellar sources (S1,2,3 and 4) are marked. The solid line represents unreddened main-sequence stars (Koornneef 1983; Bessell & Brett 1988) and the dashed line is the reddening vector of $A_V = 30$ mag. S1 shows a high extinction of $A_V = 28$. The other sources S2, S3 and S4 also show high extinctions of $A_V = 19, 26$ and 36 respectively. Based on the photometry in Table 1, us-

ing Rieke & Lebofski (1985) extinction law and a distance modulus = 11.5 mag, the expected absolute K magnitudes of the sources S1, 2, 3 and 4 are $M_K = -3.6, -2.4, -2.8$ and -3.0 respectively. Source S5 which is only observed in the K' band perhaps suffers extinction greater than $A_V = 40$ mag, if we consider that it is fainter than the detection limits of the Mt. Abu observations in J and H bands.

If the above arguments are true, then the sources have to be much more massive than T Tauri stars. Following the Palla & Stahler (1999) evolutionary tracks, stars less massive than $3.5 M_{\odot}$ (at any age) are never more luminous than $M_K = -1.67$. Assuming that more massive stars actually burn hydrogen while accreting, they are expected to have no pre-mainsequence evolution (as the authors above advocate). Using the zero-age main sequence magnitude/color relations $M_K = -3.6$ implies a spectral type between B0 and B1 and $M_K = -2.4$ corresponds to B2. We have considered IR colors from Bessell & Brett (1988) for our calculations. In all these arguments, we have neglected infrared excess and accretion luminosity.

For the source S1, being the most massive, the accretion luminosity could contribute substantially to the luminosity only if the accretion rate is very high. This is possible, but implies an amount of circumstellar material that would be so high to produce a column density much higher than $A_V = 28$ mag. For other sources S2, 3 and 4 the uncertainties in the infrared excess will apply, however, the absolute magnitude-luminosity relation does give us a lower limit for the stellar masses.

The important finding of the current photometric study is Star S5 whose J and H magnitudes are fainter than 15.2 and 16.4 magnitudes (our detection limits) while the K' magnitude is 12.9 (2MASS data gives 13.52). The absence of S5 in the H band image of HLR02 confirms this. In any case a deeper NIR survey preferably at high spatial resolution may show exactly how steep is the NIR energy distribution for S5 and hence its exact nature. The difference in the K magnitudes from Mt Abu and 2MASS is 0.62 and is larger than the observational uncertainties. It is possible that the star may be undergoing a variability that is akin to FU Orionis. Such a substantial difference is also noticed (Table 1) in the case of K band for S1.

3.2. Molecular Hydrogen Knots in IRAS 06061+2151

Figures 4 and 5 show narrow band images of the region in H_2 (2.121 μm , line + continuum) and $\text{Br}\gamma$ (2.167 μm , line + continuum) respectively. The circle marks two knot-like diffuse structures, one in the north-west and another in the south-east of the centre of the cluster in the H_2 image. These two prominent diffuse emission features are not seen in the $\text{Br}\gamma$ image. We may safely conclude then that the narrow band image at 2.167 μm (Fig 5) represents the background continuum radiation. We estimated the integrated brightness of the knots using relative photometry with respect to the Ks magnitude of S1 and S2. A large aperture of 20 arcsecs was used for the knots. We

find that in the H_2 image, the western knot is 1.6 mag and the eastern knot 2.5 mag fainter than S1. A similar estimation of brightness of faint stars seen in both the H_2 and $Br\gamma$ images showed that they are 4 mag fainter. We also find that these knot-like structures should be at least 4.5 mag fainter than S1 in the $Br\gamma$ filter since they are not detected in the image.

We thus establish that the two newly identified structures are primarily due to H_2 emission. The H_2 $v = 1-0$ S(1) $2.122\mu m$ line emission has been proven to be an excellent tracer in dense molecular cloud regions for moderate shocks in the range of 10 to 30 km/s (Shull & Beckwith 1982; Draine, Roberge, & Dalgarno 1983; Sternberg & Dalgarno 1989) and are found to trace the termination of CO outflows and optically seen HH objects (see eg. BACHILLER 1996; Hartmann 1998 and references therein). It is possible that due to this moderate nature of the shock the ionization of hydrogen could not take place. Therefore, we propose that this pair of knot-like structures is probably tracing a shock caused in the surrounding interstellar medium by protostellar jets. We call them Knot-NW and Knot-SE respectively. These knot-like structures may be considered as infrared counterparts of HH Objects that are often detected in [SII] and $H\alpha$ images. From the H_2 image, we determine the co-ordinates ($\alpha(2000)$, $\delta(2000)$) of the Knot-NW and Knot-SE as (06h09m05.2s, $+21^\circ50'56''$) and (06h09m08.5s, $+21^\circ50'32''$) respectively. Judging from the brightness of the knots, we assume that Knot-NW suffers lesser extinction and hence is tracing a blue jet and the Knot-SE the red one. Alternatively, the difference in brightness could also be due to the differences in physical parameters such as density of the two regions of interstellar medium.

There are two sources, namely S1 and S4 lying in the line of the jets. From the color-color diagram (Fig 2), it appears that S1 is likely a massive member of the cluster and a Class II source, while S4 is less massive but clearly fits in the scenario of a Class I protostar. So it is tempting to suggest that S4 could be more likely the source of the jets. It is also situated nearly in the middle of the projected distance between the two knots. The projected length of the jet is estimated to be about 0.5 pc.

3.3. Medium Resolution TNG K Band Spectra

Figure 6 shows in the upper panel normalized K band spectra, taken at TNG, towards the star S1 and in the lower panel the nebular spectra obtained in a region 5 arcsec west of S1 and towards S4 (close to the path of the NW jet; see Figure 4). Our main concern is the H_2 line ratio of 2-1 S(1)/1-0 S(1) ($2.247\mu m/2.122\mu m$), which depends critically on the type of excitation mechanism namely fluorescence or shock heating (Shull & Beckwith 1982; Draine, Roberge & Dalgarno 1983; Black & van Dishoeck 1987; Sternberg & Dalgarno 1989; Draine & Bertoldi 1996; Luhman, Engelbracht & Luhman 1998; Walmsley et al 2000; line ratios compiled in Glass 2000).

Since the $2.247\mu m$ line is very weak and buried in the base-level noise, we put an upper limit for the ratio at 0.1 and 0.05 for the nebular spectra and the Star 1 respectively by taking the base line as 1.0 ± 0.1 for nebular spectra and 1.0 ± 0.05 for the Star 1 spectra. therefore, these upper-limit ratios strongly suggest that the excitation could be due to a shock rather than fluorescence where one would expect the ratio to be ≥ 0.5 (eg., Black & van Dishoeck 1987; Draine & Bertoldi 1996). One can notice the $2.223\mu m$ line of H_2 1-0S(0) appearing quite prominently in both the spectra with the ratio 1-0S(0)/1-0S(1) equal to 0.55 for nebula and 0.35 for star 1. these ratios are rather high for collisional excitation. but at an equilibrium temperature of 1000 K, the ratio of 1-0S(0)/1-0S(1) rises (to ~ 0.3), while the ratio of 2-1S(1)/1-0S(1) falls to insignificance (see table 2 in Black & van Dishoeck 1987). The fluorescence excitation populates the vibrational levels $v\geq 2$ much more than in the thermal excitation; this does not seem to be the case here. We therefore infer from the spectra that the lines are likely thermally excited. The ratios for S1 are in reasonably good agreement with the ratios obtained by HLR02. Prominently missing in the spectra is the $Br\gamma$ emission line at $2.167\mu m$ which could be due to the fact that the line if present is very weak. The absence of $Br\gamma$ is also reported by HLR02. So it is clear from the presented spectra that the molecular hydrogen excitation is caused by moderate shocks (with velocities less than about 30 km/s that can not cause hydrogen ionization as it requires greater than 60 km/s) which explains why $Br\gamma$ emission line is absent or very weak.

In Fig 6 we find the spectrum towards S1 (upper panel) to be significantly weaker than the spectrum on the nebular region towards S4 (lower panel). So if S1 is indeed a class I protostar then we expect the two spectra to be of more or less comparable strengths (since S4 is a protostar). This is a further evidence that S1 is likely a massive and more evolved member of the cluster.

HLR02 concluded that the radio source KCW 188.793+1.030 (R1) corresponds to the source S1 while the spectra taken by them on the radio source KCW 188.796+1.030 (R2) do not seem to have a counterpart in their H band image. But our K' image taken at Mt Abu as well as the TNG narrow band images show that the S5 source does line up with HLR02 spectra on R2. The line ratios reported by HLR02 on R2 (S5) are nearly similar to those on R1 (S1). The absence of $Br\gamma$ in the two spectra presented by HLR02 and the TNG spectra shows that in general the region may be pervaded by a mild shock with velocities not greater than 30km/s.

4. Conclusion

IRAS 06061+2151 appears to be a cluster of at least 5 bright sources four of which seem to be early B type YSOs. From the J-H/H-Ks color-color diagram we found that one of them, S1, could be the most massive member of the cluster rather than a background source and S4 is a Class I type. S3 seems to be a Class II source while S2 is

likely to be Class I type. The IRAS source is also associated with a nebulosity. As many as 3-4 Class I (including S4 and maybe S2) and 13-14 Class II/III sources (including S1 and S3) were found towards the region of IRAS 06061+2151 from the color-color diagram which confirms the earlier studies that the region contains large number of YSOs. A new source S5 appears to have a very steep infrared energy distribution and is likely a massive protostar as its colors (lower limits) suggest a heavily embedded source.

From the narrow band images in H_2 and $Br\gamma$, two oppositely directed knot-like structures (Knot-NW and Knot-SE) have been detected. The knots are likely generated by jets from the protostellar source S4 and are the infrared counterparts of the classical HH objects in optical region. From the ratios of H_2 lines, we conclude that the region is probably pervaded by a mild shock or a photodissociation region that can excite these lines but not strong enough to excite the HI lines.

Acknowledgements. We thank the referee Dr. M. Hoogerheijde for valuable suggestions for improving the original version. It is a pleasure to thank the staff at Mt. Abu IR Telescope and the TNG, La Palma facility for excellent support during observations. This work is based in part on the observations made with the Italian Telescopio Nazionale Galileo at La Palma operated by the Centro Galileo Galilei of CNAA at the Observatorio del Roque de los Muchachos of the IAC, Canary Islands, Spain. This publication makes use of data products from the 2MASS, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center, funded by the NASA and the NSF. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

References

- Anandarao, B.G., 1998, In High Energy Astrophysics, Univ. Press, Hyderabad
- Bachiller, R., 1996, ARA&A, 34, 111
- Baffa, C., et al. 2001, A&A, 378, 722
- Bessel, M., & Brett, J.M., 1988, PASP, 100, 1134
- Bica, E., Dutra, C.M., & Barbuy, B., 2003, A&A, 397, 177
- Black, J.H., & van Dishoeck, E.F., 1987, ApJ, 322, 412
- Carpenter, J.M., et al. 1993, ApJ, 407, 657
- Carpenter, J.M., Snell, R.L., & Schloerb, F.P., 1995a, ApJ, 445, 246
- Carpenter, J.M., Snell, R.L., & Schloerb, F.P., 1995b, ApJ, 450, 201
- Chakraborty, A., Ojha, D.K., Anandarao, B.G., & Renagarajan, T.N., 2000, A&A, 364, 683
- Davis, C.J., Eisloffel, J., & Ray, T.P., 1994, ApJ, 426, L93
- Draine, B.T., & Bertoldi, F., 1996, ApJ, 468, 269
- Draine, B.T., Roberge, W.G., & Dalgarno, A., 1983, ApJ, 264, 485
- Eisloffel, J., Smith, M.D., & Davis, C.J., 2000, A&A, 359, 1147
- Ghosh, S.K., Iyengar, K.V.K., Karnik, A.D., Rengarajan, T.N., et al. 2000, BASI, 28, 515
- Glass, I.S., Handbook of Infrared Astronomy, Cambridge Univ. Press, Cambridge, 1999
- Gomez, M., Kenyon, S.J., & Hartmann, L., 1994, AJ, 105, 1850
- Hanson, M.M., Luhman, K.L., & Rieke, G.H., 2002, ApJSS, 138, 35
- Hartmann, L., Accretion Processes in Star Formation, Cambridge Univ. Press, Cambridge, 1998
- Herbst, T.M., Beckwith, S., & Robberto, M., 1997, ApJ, 486, L59
- Hunter, T.R., Testi, L., Taylor, G.B., et al. 1995, A&A, 302, 249
- Kenyon, S.J., Whitney, B.A., Gomez, M., et al. 1993, ApJ, 414, 773
- Koornneef, J., 1983, A&A, 128, 84
- Kurtz, S.E., Churchwell, E., & Wood, D.O.S., 1994, ApJSS, 91, 659
- Lada, C.J., & Adams, F.C., 1992, ApJ, 393, 278 L
- Lada, C.J., Young, E.T., & Greene, T.P., 1993, ApJ, 408, 471 L
- Luhman, K.L., Engelbracht, C.W., & Luhman, M.L., 1998, ApJ, 499, 799
- Meyer, M.R., Calvet, N., & Hillenbrand, L.A., 1997, AJ, 114, 288
- Nandakumar, M.S., PhD Thesis, Gujarat University, Ahmedabad, 1999
- Palla, F., & Stahler, S.W., 1999, ApJ, 525, 772
- Rieke, G.H., & Labofsky, M.J., 1985, ApJ, 288, 618
- Schwartz, R.D., Williams, P.M., Cohen, M., & Jennings, D.G., 1988, ApJ, 334, 99
- Shepherd, D.S., & Churchwell, E., 1996, ApJ, 457, 267
- Shull, J.M., & Beckwith, S., 1982, ARAA, 20, 163
- Sternberg, A., & Dalgarno, A., 1989, ApJ, 338, 197
- Stetson, P.B., 1987, PASP, 99, 191
- Strom, K.M., Margulis, M., & Strom, S.E., 1989, ApJ, 346, L33
- Walmsley, C.M., Natta, A., Oliva, E., & Testi, L., 2000, A&A, 364, 301




Fig. 1. K' band image of IRAS 06061+2151 observed from Mt Abu. The numbers represent the cluster of sources around IRAS 06061+2151 region. The abscissa and ordinate are in J2000.0 epoch.




Fig. 2. Color-color diagram of the sources extracted from the 2MASS-PSC data. The numbers represent the cluster of sources around IRAS 06061+2151 region (see text and Table 1 for details). The solid line represents unreddened main-sequence stars and the giants (Bessell & Brett 1988). The dashed lines are parallel to the reddening vector with magnitude of $A_v=30$. The dotted line is the locus of CTTS (Meyer et al. 1997).




Fig. 3. Color-Magnitude diagram of the sources extracted from the 2MASS-PSC data. The numbers represent the cluster of sources around IRAS 06061+2151 region. The solid line represents unreddened main-sequence stars (Koornneef 1983; Bessell & Brett 1988) and the dashed line is the reddening vector with magnitude of $A_v=30$.






Fig. 4. H_2 line + continuum (2.122 μm) image observed from TNG. The circles show the Knot-like objects Knot-NW and Knot-SE. The numbers 1,2,3,4 and 5 mark the positions of the stars as numbered in Table 1. North is up and East is to the left. The abscissa and ordinate are in J2000.0 epoch.

Table 1. A comparison of stellar magnitudes from 2MASS data and Mt. Abu images.

No.	Co-ordinates (J2000) RA-DEC	2MASS			MtAbu		
		<i>J</i>	<i>H</i>	<i>K_s</i>	<i>J</i>	<i>H</i>	<i>K'</i>
1	06h09m07.2s +21°50'40''	15.46	12.87	11.22	15.54	12.68	10.85
2	06h09m07.1s +21°50'34''	13.67	12.14	11.20	13.65	12.24	11.11
3	06h09m07.5s +21°50'32''	15.31	12.94	11.55	15.44	13.05	11.45
4	06h09m06.9s +21°50'43''	—	14.34	12.16	16.66	14.48	12.29
5	06h09m07.1s +21°50'32''	—	—	13.52	—	—	12.90

**Fig. 5.** Br γ line + continuum (2.167 μ m) image observed from TNG. North is up and East is to the left. The abscissa and ordinate are in J2000.0 epoch.**Fig. 6.** Spectra towards Star 1(top) and that of nebula (bottom) which is 5 arcsec west of Star 1 towards Star 4. Both the spectra are normalized with respect to its continuum.